

# Paleomagnetic evidence for a Tertiary not Triassic age for rocks in the lower part of the Grober-Fuqua #1 well, southeastern Albuquerque Basin, New Mexico

Mark R. Hudson and V. J. S. Grauch, U.S. Geological Survey, Denver, Colorado

## Abstract

A sedimentary sequence penetrated in the lower part of the Grober-Fuqua #1 well in the southeastern Albuquerque Basin has previously been interpreted as either Triassic or Eocene in age. Paleomagnetic study of three specimens from two core fragments yielded a 54.5° mean inclination of remanent magnetization relative to bedding. This inclination is like that expected in Tertiary time and is distinct from an expected low-angle Triassic inclination. Although the data are very few, when considered in combination with stratigraphic relations and the presence of a gravity low in this southeastern part of the basin, the paleomagnetic evidence favors a Tertiary

age for strata in the lower part of the Grober-Fuqua #1 well.

## Introduction

The Albuquerque Basin is part of the Rio Grande rift, which extends from Colorado to Mexico. Basin subsidence was most active from late Oligocene through Miocene time (Chapin and Cather, 1994), resulting in large accumulations of syn-rift sedimentary deposits of the Santa Fe Group. The Santa Fe Group varies in thickness from approximately 1,000 to 2,000 m (3,280 to 6,560 ft) along the basin margins

to more than 4,270 m (14,000 ft) in the basin centers (Lozinsky, 1994; May and Russell, 1994). Throughout most of the Albuquerque Basin, the thickest sections of Santa Fe Group correspond to areas of deep lows on gravity maps (Cordell, 1976; Birch, 1982). An apparent exception is located in the southeastern part of the Albuquerque Basin east of the Rio Grande (eastern part of the Belen sub-basin), where an extensive, north-south elongated, >20-mGal gravity low was intersected by the 1,920-m- (6,300-ft-) deep Grober-Fuqua #1 petroleum exploration well (Fig. 1). Interpretations of well data (Fig. 2) indi-





FIGURE 1—Generalized geology of the southern Albuquerque Basin from Kelley (1977), showing locations of oil exploration wells completed to >1,000 m (3,280 ft) total depth (TD; Black, 1982, 1999). GF1 = Grober-Fuqua

#1 well. Isostatic residual gravity from Gillespie et al. (2000). Contour interval = 6 milligals. HB = Hubbell Bench, where Triassic rocks were sampled by Lozinsky (1988).



\*sampled for this study

FIGURE 2—Schematic diagram of lithology, interpreted picks, and depths of core samples from the Grober-Fuqua #1 well. Interpreted picks are from R. L. Bates, as reported in Reiche (1949, p. 1204), Foster (1978), Lozinsky (1988), and Brister (written comm. 2003). The two completion depths (TD) from 1940 and 1946 and the limited depth extent of the electric log are also shown. Lithologic descriptions are from Foster (1978 and unpublished

notes) and Lozinsky (1988). Horizons where core samples are available in the lower part of the well and those (GC samples) used in Lozinsky's (1988, 1994) analysis of sandstone petrology are marked with thick black ticks. One specimen from GC-4 and two specimens from GP-1 were used in this study.

cate the Santa Fe Group is 1,384 m (4,540 ft) thick (Foster, 1978; Lozinsky, 1988, 1994). This thickness contradicts gravity models that indicate a Santa Fe Group thickness considerably greater than 2,000 m (6,560 ft; Cordell, 1976; Birch, 1982; Grauch et al., 1999). Coverage of published seismic data is minimal in this area (Russell and Snelson, 1994).

One way to reconcile the apparent con-

tradiction is by altering the gravity models to provide for low-density salt within units underlying the Santa Fe Group, as argued by Grauch et al. (1999). However, estimating the amount of salt and speculating on its geologic significance hinge on correct identification of the older units penetrated by the well. Cuttings from below 1,384 m (4,540 ft) in the Grober-Fuqua #1 well have been identified in the literature alternative-

ly as pre-rift sedimentary deposits of early Tertiary age (Baca Formation?, Foster, 1978) or Triassic clastic sedimentary rocks (Lozinsky, 1988). If the units are Triassic in age, gravity models would require an extensive amount of salt, which most likely was deposited during the Permian. For example, a preferred gravity model of Grauch et al. (1999) incorporated salt as 30% of a 2–3-km- (1.2–1.8-mi-) thick Paleo-

zoic basin-fill sequence. If the units are early Tertiary in age instead, gravity models would not require as much salt, and the salt may have been deposited irregularly in syn-basin playa lakes. Thus, a correct age determination of these units is important for understanding the significance and tectonic implications of the gravity low.

In an attempt to resolve the age controversy, we have applied paleomagnetic methods to two core fragments taken from the lower part of the Grober-Fuqua #1 well. Paleomagnetic study of these samples can discriminate between the two ages because directions of geomagnetic fields at the well site were much different in Tertiary time than in Triassic time. This difference is well expressed in the expected inclination of the field, which was 50°–60° for Tertiary time and less than 10° for Triassic time (Fig. 3). Although the two core samples from the Grober-Fuqua #1 well are unoriented, bedding is recognized. Inclination of remanent magnetization is determined relative to this paleohorizontal marker.

### Lithology of the Grober-Fuqua #1 well

A drilling summary (scout card), drillers' logs, several lithologic logs, a geophysical electric log (limited in depth), well cuttings, and core fragments from the Grober-Fuqua #1 well are available at the New Mexico Bureau of Geology and Mineral Resources (NMBGMR) petroleum library. The well was drilled to a total depth of 1,212 m (3,975 ft) in 1940. In 1946 it was redrilled to a depth of 1,920 m (6,300 ft). Cuttings and core at various intervals were collected below 802 m (2,630 ft) during drilling. Small, bagged samples from the cuttings and fragments of some of the core remain at NMBGMR.

Previous workers collectively have identified the first 1,384 m (4,540 ft) of sediments in the well as Quaternary alluvium and Santa Fe Group (Fig. 2). Interpretation on the interval below 1,384 m (4,540 ft) has changed with time (Fig. 2). This interval has been interpreted to represent Upper Cretaceous and Triassic rocks (R. L. Bates, reported in Reiche, 1949, p. 1204); pre-Santa Fe Tertiary units, possibly equivalent to the Baca Formation of Eocene age (Foster, 1978); and entirely Triassic rocks (Lozinsky, 1988).

The Triassic interpretation by Lozinsky (1988) came out of his study of sandstone petrology from cores and surface rocks from throughout the Albuquerque Basin. He observed that sandstone from core sample GC-4 (Fig. 2) differed significantly from the sandstone petrology of the Santa Fe Group, the Baca Formation, and other pre-Santa Fe Tertiary units that he had observed. Instead, petrology of GC-4 matched that of a sandstone sample from




FIGURE 3—Plot of expected inclination versus age and inclination results for the Grober-Fuqua #1 well. Inclinations are calculated from poles of Besse and Courtillot (1991) for ages of 10 Ma to 180 Ma and from the North American craton poles of Molina-Garza et al. (1998) for Early Jurassic through Triassic ages. Also shown is the mean 54.5° inclination for the Grober-Fuqua #1 well samples and the 95% confidence limits (shaded) for its comparison to the expected inclinations.

Triassic rocks mapped by Kelley (1977) at the Hubbell Bench (Fig. 1) that Lucas (1991) correlated with basal units of the Triassic section. The lithologic descriptions of the lower part of the well also resemble descriptions of outcrops of Triassic rocks on the flanks of the basin (Stewart et al., 1972; Lucas and Hayden, 1989; Lucas and Heckert, 1994, 1995). Notable Triassic lithologic similarities include abundant mica, bentonitic clays, subangular to subrounded grains, and the presence of conglomerates.

### Paleomagnetism

Inspection of available samples from the Grober-Fuqua #1 well revealed that only core fragments from the GC-4 and GP-1 horizons (Fig. 2) were both sufficiently large for rock magnetic analysis and had visible bedding laminations. Roughly cube-shaped specimens of approximately 1 cc volume were cut from these core fragments and oriented in specimen coordinates so that bedding was horizontal. The GC-4 specimen, from the 1,541–1,544 m (5,057–5,064 ft) interval, is pale reddish-brown (10R 5/4), laminated, very fine to fine-grained sandstone. Two specimens were obtained from GP-1, from the 1,837–1,838 m (6,026–6,031 ft) interval; they are pale-red (10R 6/2), very fine to medium-grained sandstone with laminations of grayish-red (10R 4/2) shale.

The specimens were subjected to progressive thermal and alternating-field

demagnetization to isolate characteristic magnetization components and assess their origin. Remanent magnetization of the specimens was measured using a spinner magnetometer with a sensitivity of about 1E-4 A/m; two measurements were made and then averaged for each specimen at each demagnetization step. Thermal demagnetization of the GC-4 specimen isolated one component of magnetization above 395° C with an unblocking range extending as high as 675° C (Fig. 4A). Principal component analysis (Kirschvink, 1980) of these demagnetization data yielded an inclination of 58.2° for this remanent magnetization component with an associated maximum angular deviation (MAD) of 4.2°. Thermal demagnetization behavior for the GP-1 specimen was more complex. A moderately inclined component of remanent magnetization (61.1°, MAD of 8.2°) was isolated over 350°–560° C (Fig. 4B), but at higher temperatures the directions and magnitudes of magnetization varied in an erratic manner and no coherent magnetization component was isolated. Alternating-field demagnetization of the second specimen from the GP-1 horizon isolated a magnetization of moderate inclination (44.3°, MAD of 7.5°) over peak inductions of 25 mT to 100 mT (Fig. 4C).

For GC-4 the unblocking temperatures as high as 675° C indicate a hematite carrier for the characteristic component, whereas for the GP-1 specimens the removal of the characteristic component over ranges




FIGURE 4—Orthogonal vector diagrams (Zijderveld, 1967) for progressive demagnetization results of the GC-4 and GP-1 specimens. Solid symbols are projections of remanent magnetizations after each demagnetization step on the horizontal plane, and they represent arbitrary declinations. Open symbols are projections of remanent magnetizations after each

demagnetization step on the vertical plane, and they represent inclinations relative to horizontal. Values for GC-4 and GP-1 #1 are demagnetization temperatures ( $^{\circ}\text{C}$ ); values for GP-1 #2 are peak inductions (mT). Dashed lines are mean inclinations determined over the selected demagnetization ranges for the characteristic magnetization component of each specimen.

of  $300^{\circ}$ – $560^{\circ}\text{C}$  and  $25$ – $100\text{ mT}$  indicates a low-Ti titanomagnetite carrier. These conclusions were supported by isothermal remanent magnetization (IRM) acquisition experiments (Fig. 5). In IRM experiments, titanomagnetite acquires large magnetizations at inductions as high as  $0.3\text{ T}$ , but at higher inductions it becomes saturated and acquires no additional IRM (Dunlop, 1986). The GP-1 specimens acquired at least 90% of their IRM by inductions of  $0.3\text{ T}$ , consistent with a main titanomagnetite carrier (Fig. 5). In contrast, the GC-4 specimen acquired 64% of the IRM at inductions above  $0.3\text{ T}$ , requiring a high-coercivity carrier like hematite.

Based on statistical procedures of McFadden and Reid (1982), the mean inclination of the three specimens is  $54.5^{\circ} \pm 32.4^{\circ}$ . We include both specimens from GP-1 in the mean calculation because their inclination range is greater than their difference with the GC-4 inclination. The associated 95% confidence limit for the mean inclination is large due to the small number of samples available. To assess the age of the magnetization we compared the mean inclination for the Grober-Fuqua #1 well samples to expected inclinations at the site calculated from North American reference poles. For Triassic through Early Jurassic times we used the North American craton poles of Molina-Garza et al. (1998) for reference. For Late Jurassic and younger times we used the North American poles from Besse and Courtillot (1991). Uncertainty limits for the inclination differences are  $24^{\circ}$ – $25^{\circ}$ , calculated using the

flattening parameter of Demarest (1983). Thus the  $54.5^{\circ}$  mean inclination for the Grober-Fuqua #1 well samples is similar to values expected for Tertiary and Cretaceous times, and it is statistically distinct from expected Triassic inclinations (Fig. 3).

Interpretation of the inclination data to indicate a Tertiary age for the sedimentary rocks in the lower Grober-Fuqua #1 well requires two principal assumptions: (1) that bedding preserved in the core samples was horizontal when the rocks acquired their remanent magnetization, and (2) that the age of the remanent magnetization approximates the age of the sedimentary sequence. Hypothetically, a properly oriented primary dip of bedding (e.g., cross-bedding) could cause a low-inclination Triassic magnetization to appear to have a steeper inclination when the bedding is oriented horizontally. To produce the  $54.5^{\circ}$  inclination, however, the primary dip from both well horizons must have been high ( $45^{\circ}$ – $50^{\circ}$ ) and directed opposite to a recorded Triassic paleofield declination (generally north-northwest or south-southeast). These circumstances seem unlikely.

If the moderately inclined remanent magnetization measured in the well resulted from remagnetization of Triassic rocks, this remagnetization event was likely recorded in other Triassic rocks in the region. Remanent magnetizations in nearby Triassic sections were examined by Molina-Garza et al. (1991) and Molina-Garza and Geissman (1999). Molina-Garza et al. (1991) isolated low-inclination Trias-

sic magnetizations carried by hematite in most samples of Moenkopi and Chinle Formations from rift-flank uplifts to the northeast, west, and south of the Grober-Fuqua #1 well. For example, from their Tables 2 and 3, they included results from 281 of 351 demagnetized samples (79%) from 53 sites used in their calculation of Triassic mean directions. The Triassic magnetization component was typically unblocked over  $580^{\circ}$ – $680^{\circ}\text{C}$ . Secondary magnetizations of moderate to steep inclination also were recognized at several locations, and these were concentrated above the basal disconformity with Permian strata (Molina-Garza and Geissman, 1999). The secondary magnetizations are either pretilting reversed-polarity components or post-tilting normal polarity components, and they are carried by goethite or fine-grained hematite. These secondary components typically unblocked over lower and more distributed temperature ranges than the primary low-inclination components.

For the Grober-Fuqua #1 well samples, the finding that magnetite carries the characteristic magnetization in the GP-1 horizon is an important consideration. The reddish color of this sample suggests that it was never subjected to a reducing geochemical environment that could precipitate authigenic magnetite. Thus the magnetite in GP-1 is most likely detrital in origin, and it probably acquired a detrital or post-detrital magnetization at or soon after sediment deposition. Later thermoviscous remagnetization of this early-formed mag-

netite remanence seems unlikely because it would require a thermal pulse of approximately 525° C for 10 m.y. to remagnetize the magnetite that unblocks as high as 560° C in laboratory demagnetization (Pulliah et al., 1975). For GC-4, the remanent magnetization carried by hematite is probably a chemical remanent magnetization formed after sediment deposition. Its unblocking as high as 675° C is more like that expected for coarse specular hematite rather than for fine-grained hematite that Molina-Garza et al. (1991) interpreted to typically carry secondary magnetizations in Triassic rocks. The hematite remanence in GC-4 may well have formed within a few million years after deposition, as is common in red bed sequences (Butler, 1992).

## Discussion

The paleomagnetic results support a Tertiary age for rocks penetrated in the lower part of the Grober-Fuqua #1 well, but we interpret the data with caution because they are based on so few samples. However, in combination with several regional geologic considerations, the results argue against a Triassic age. First, the depth range of the lower interval in the well would require a Triassic section of more than 536 m (1,760 ft) thick (Fig. 2). This is unreasonable considering the maximum thickness of the Triassic section thins from 480–500 m (1,575–1,640 ft) on the northern and northeastern flanks of the Albuquerque Basin (Lucas and Heckert, 1995; Lucas et al., 2001) to less than 270 m (886 ft) in south-central New Mexico (Lucas, 1991). Second, Jurassic or Cretaceous sandstones typically overlie Triassic rocks from the western to the southeastern sides of the Albuquerque Basin (Lucas and Heckert, 1994; Lucas, 1991). The absence of these distinctive units in the well unreasonably implies that Tertiary units rest directly on Triassic rocks (S. Lucas, written comm. 2003). Finally, the Triassic pick in the well requires surprising explanations for the presence of the extensive gravity low in the area (Fig. 1), leading inevitably to the conclusion that a significant volume of low-density salt exists within the normally dense Mesozoic–Paleozoic sedimentary sequence (Grauch et al., 1999). Although the presence of Permian salt is not unreasonable, large quantities probably would have mobilized into diapirs through time, particularly during Laramide shortening. No evidence of salt tectonism is known in the region.

Choosing among the previously published interpretations for the lower part of the Grober-Fuqua #1 well (Fig. 2), several arguments favor Foster's (1978) interpretation that the strata are correlative with the Eocene Baca Formation. First, it is reasonable to expect that Baca Formation associated with the Eocene Carthage–La Joya

Basin, exposed as near as 40 km (25 mi) to the south in the Joyita Hills, would extend northward into the area of the Grober-Fuqua #1 well (Cather and Johnson, 1984; Cather et al., 1994). Second, the Manzano Mountains (Fig. 1) were high and undergoing denudation during Eocene time, as was the Hubbell Bench but to a lesser degree (Kelley et al., 1992). These areas could have served as nearby sources of Triassic-rock detritus.

On the other hand, if the lithology of the lower part of the Grober-Fuqua #1 well is no longer considered to be diagnostic of a Triassic or an early Tertiary age, this interval could be a continuation of the Santa Fe Group of Miocene or late Oligocene age. In a recent reexamination of the well cuttings and core fragments, B. Brister (written comm. 2003) has favored a Santa Fe Group identity for the lower well sequence (Fig. 2) based on an assemblage of lithic fragments that includes plutonic and metamorphic types of probable Precambrian provenance and basaltic and ash-flow tuff types of likely Oligocene or Miocene age. This hypothesis is also consistent with the paleomagnetic data (Fig. 3), and it would be favored by previous gravity inversions that inferred rift-fill thickness in excess of 2,000 m (6,560 ft) in the southeastern part of the Albuquerque Basin (Cordell, 1976; Birch, 1982; Grauch et al., 1999).

## Conclusions

Although the available samples are few, inclinations of remanent magnetization determined from the lower part of the Grober-Fuqua #1 well are like those expected for Tertiary time, and they are distinct from those expected for Triassic time. Coupled with the recognition that the penetrated stratigraphic section in the lower part of the well is abnormally thick




FIGURE 5—Acquisition of isothermal remanent magnetization of Grober-Fuqua #1 well core samples. Backfield demagnetization of remanence acquired at 1.2 T gives coercivity of remanence values that are indicative of hematite for the GC-4 horizon (triangles) and magnetite for the GP-1 horizon (squares and diamonds).

for an expected Triassic section and also with the presence of a gravity low in this southeastern part of the basin, the paleomagnetic data provide support for a Tertiary age for sandstones in the lower part of the Grober-Fuqua well.

## Acknowledgments

We thank Ron Broadhead of the NMBGMR for kindly providing samples from the Grober-Fuqua #1 well. The second author greatly appreciates the many stimulating discussions provided by geologists on both sides of "the Grober problem." Bruce Black, Ron Broadhead, Dave Love, Brian Brister, and Rick Lozinsky were especially generous with their time and energy. We appreciate reviews by Brian Brister, Florian Maldonado, and Richard Reynolds.

## References

- Besse, J., and Courtillot, V., 1991, Revised and synthetic apparent polar wander paths of the African, Eurasian, North American, and Indian plates, and true polar wander since 200 Ma: *Journal of Geophysical Research*, v. 96, pp. 4029–4050.
- Birch, F. S., 1982, Gravity models of the Albuquerque Basin, Rio Grande rift, New Mexico: *Geophysics*, v. 47, no. 8, pp. 1185–1197.
- Black, B. A., 1982, Oil and gas exploration in the Albuquerque Basin; in Grambling, J. A., Wells, S. G., and Callender, J. F. (eds.), *Albuquerque country II: New Mexico Geological Society, Guidebook 33*, pp. 313–324.
- Black, B. A., 1999, Recent oil and gas exploration in the Albuquerque Basin; in Pazzaglia, F. J., and Lucas, S. G. (eds.), *Albuquerque geology: New Mexico Geological Society, Guidebook 50*, pp. 437–440.
- Butler, R. F., 1992, *Paleomagnetism—magnetic domains to geologic terranes*: Blackwell Scientific, Boston, 319 pp.
- Cather, S. M., Chamberlin, R. M., and Ratté, J. C., 1994, Tertiary stratigraphy and nomenclature for western New Mexico and eastern Arizona; in Chamberlin, R. M., Kues, B. S., Cather, S. M.,

- Barker, J. M., and McIntosh, W. C. (eds.), Mogollon slope, west-central New Mexico and east-central Arizona: New Mexico Geological Society, Guidebook 45, pp. 259–266.
- Cather, S. M., and Johnson, B. D., 1984, Eocene tectonics and depositional setting of west-central New Mexico and eastern Arizona: New Mexico Bureau of Mines and Mineral Resources, Circular 192, 33 pp.
- Chapin, C. E., and Cather, S. M., 1994, Tectonic setting of the axial basins of the northern and central Rio Grande rift; *in* Keller, G. R., and Cather, S. M. (eds.), Basins of the Rio Grande rift—structure, stratigraphy, and tectonic setting: Geological Society of America, Special Paper 291, pp. 5–23.
- Cordell, L., 1976, Aeromagnetic and gravity studies of the Rio Grande graben in New Mexico between Belen and Pilar; *in* Woodward, L. A., and Northrop, S. A. (eds.), Tectonics and mineral resources of southwestern North America: New Mexico Geological Society, Special Publication 6, pp. 62–70.
- Demarest, H. H., 1983, Error analysis of the determination of tectonic rotation from paleomagnetic data: *Journal of Geophysical Research*, v. 88, pp. 4321–4328.
- Dunlop, D. J., 1986, Coercive forces and coercivity spectra of submicron magnetites: *Earth and Planetary Science Letters*, v. 78, pp. 288–295.
- Foster, R. W., 1978, Selected data for deep drill holes along Rio Grande rift in New Mexico; *in* Hawley, J. W. (comp.), Guidebook to Rio Grande rift in New Mexico and Colorado: New Mexico Bureau of Mines and Mineral Resources, Circular 163, pp. 236–237.
- Gillespie, C. L., Grauch, V. J. S., Oshetski, K., and Keller, G. R., 2000, Principal facts for gravity data collected in the southern Albuquerque Basin and a regional compilation, central New Mexico: U.S. Geological Survey, Open-file Report 00-490, digital report available at <http://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-00-240>.
- Grauch, V. J. S., Gillespie, C. L., and Keller, G. R., 1999, Discussion of new gravity maps for the Albuquerque Basin area; *in* Pazzaglia, F. J., and Lucas, S. G. (eds.), Albuquerque geology: New Mexico Geological Society, Guidebook 50, pp. 119–124.
- Kelley, S. A., Chapin, C. E., and Corrigan, J., 1992, Late Mesozoic to Cenozoic cooling histories of the flanks of the northern and central Rio Grande rift, Colorado and New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 145, 39 pp.
- Kelley, V. C., 1977, Geology of Albuquerque Basin, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 33, 59 pp.
- Kirschvink, J. L., 1980, The least-square line and plane and the analysis of paleomagnetic data: *Geophysical Journal of the Royal Astronomical Society*, v. 62, pp. 699–718.
- Lozinsky, R. P., 1988, Stratigraphy, sedimentology, and sand petrology of the Santa Fe Group and pre-Santa Fe Tertiary deposits in the Albuquerque Basin, central New Mexico: Unpublished Ph.D. dissertation, New Mexico Institute of Mining and Technology, Socorro, 298 pp.
- Lozinsky, R. P., 1994, Cenozoic stratigraphy, sandstone petrology, and depositional history of the Albuquerque Basin, central New Mexico; *in* Keller, G. R., and Cather, S. M. (eds.), Basins of the Rio Grande rift—structure, stratigraphy, and tectonic setting: Geological Society of America, Special Paper 291, pp. 73–82.
- Lucas, S. G., 1991, Triassic stratigraphy, paleontology and correlation, south-central New Mexico; *in* Barker, J. M., Kues, B. S., Austin, G. S., and Lucas, S. G. (eds.), Geology of the Sierra Blanca, Sacramento, and Capitan Ranges, New Mexico: New Mexico Geological Society, Guidebook 42, pp. 243–259.
- Lucas, S. G., and Hayden, S. N., 1989, Triassic stratigraphy of west-central New Mexico; *in* Anderson, O. J., Lucas, S. G., Love, D. W., and Cather, S. M. (eds.), Southeastern Colorado Plateau: New Mexico Geological Society, Guidebook 40, pp. 191–211.
- Lucas, S. G., and Heckert, A. B., 1994, Triassic stratigraphy in the Lucero uplift, Cibola, Valencia, and Socorro Counties, New Mexico; *in* Chamberlain, R. M., Kues, B. S., Cather, S. M., Barker, J. M., and McIntosh, W. C. (eds.), Mogollon slope, west-central New Mexico and east-central Arizona: New Mexico Geological Society, Guidebook 45, pp. 241–254.
- Lucas, S. G., and Heckert, A. B., 1995, Triassic stratigraphy around the Sandia uplift, central New Mexico; *in* Bauer, P. W., Kues, B. S., Dunbar, N. W., Karlstrom, K. E., and Harrison, B. (eds.), Geology of the Santa Fe region, New Mexico: New Mexico Geological Society, Guidebook 46, pp. 233–241.
- Lucas, S. G., Heckert, A. B., and Hunt, A. P., 2001, Triassic stratigraphy, biostratigraphy and correlation in east-central New Mexico; *in* Lucas, S. G., and Ulmer-Scholle, D. S. (eds.), Geology of the Llano Estacado: New Mexico Geological Society, Guidebook 52, pp. 85–102.
- May, S. J., and Russell, L. R., 1994, Thickness of the syn-rift Santa Fe Group in the Albuquerque Basin and its relation to structural style; *in* Keller, G. R., and Cather, S. M., (eds.), Basins of the Rio Grande rift—structure, stratigraphy, and tectonic setting: Geological Society of America, Special Paper 291, pp. 113–123.
- McFadden, P. L., and Reid, A. B., 1982, Analysis of palaeomagnetic inclination data: *Geophysical Journal of the Royal Astronomical Society*, v. 69, pp. 307–319.
- Molina-Garza, R. S., Acton, G. D., and Geissman, J. W., 1998, Carboniferous through Jurassic data and their bearing on rotation of the Colorado Plateau: *Journal of Geophysical Research*, v. 1003, pp. 24,179–24,188.
- Molina-Garza, R. S., and Geissman, J. W., 1999, Remagnetization along the Permian–Triassic disconformity in central New Mexico and remanence acquisition in the Moenkopi Formation; *in* Pazzaglia, F. J., and Lucas, S. G. (eds.), Albuquerque geology: New Mexico Geological Society, Guidebook 50, pp. 125–132.
- Molina-Garza, R. S., Geissman, J. W., van der Voo, R., Lucas, S. G., and Hayden, S. N., 1991, Paleomagnetism of the Moenkopi and Chinle Formations in central New Mexico—implications for polar wander paths and Triassic magnetostratigraphy: *Journal of Geophysical Research*, v. 96, pp. 14,239–14,262.
- Pulliah, G. E., Irving, E., Buchan, K. L., and Dunlop, D., 1975, Magnetization changed caused by burial and uplift: *Earth and Planetary Science Letters*, v. 28, pp. 133–143.
- Reiche, P., 1949, Geology of the Manzanita and north Manzano Mountains, New Mexico: Geological Society of America, Bulletin, v. 60, no. 7, pp. 1183–1212.
- Russell, L. R., and Snelson, S., 1994, Structure and tectonics of the Albuquerque Basin segment of the Rio Grande rift—insights from reflection seismic data; *in* Keller, G. R., and Cather, S. M. (eds.), Basins of the Rio Grande rift—structure, stratigraphy, and tectonic setting: Geological Society of America, Special Paper 291, pp. 83–112.
- Stewart, J. H., Poole, F. G., and Wilson, R. F., 1972, Stratigraphy and origin of the Chinle Formation and related Upper Triassic strata in the Colorado Plateau region: U.S. Geological Survey, Professional Paper 690, 336 pp.
- Zijderveld, J. E. A., 1967, AC demagnetization of rocks—analysis of results; *in* Collinson, D. W., Creer, K. M., and Runcorn, S. K. (eds.), Methods in paleomagnetism: Elsevier, Amsterdam, pp. 254–286.