

- Circular 2, 15 pp.
- Sherer, R. L., 1990, Pajarito yttrium-zirconium deposit, Otero County, New Mexico: *New Mexico Geology*, v. 12, p. 21.
- Singer, D. A., 2000, Descriptive model of carbonatite deposits; in USGS Mineral Deposit Models: U.S. Geological Survey, Digital Data Series DDS-064, model 10, 3 pp.
- Staatz, M. H., 2000, Descriptive model of thorium-rare-earth veins; in USGS Mineral Deposit Models: U.S. Geological Survey, Digital Data Series DDS-064, model 11d, 6 pp.
- Taylor, S. R., and McLennan, S.M., 1985, *The Continental Crust; its composition and evolution*: Blackwell Science Publishers, Oxford, 312 pp.
- USGS, 2015, Mineral commodity summaries: U.S. Geological Survey, 196 pp.
- Vance, Z., 2013, *Mineralogy, Geochemistry and Genesis of the Hydrothermal REE-Fluorite-AG-PB-CU Ore Deposits of the Gallinas Mountains, New Mexico [M.S. thesis]*: Socorro, New Mexico Institute of Mining and Technology, 219 pp.
- Verplanck, P.L., Van Gosen, B.S., Seal, R.R., and McCafferty, A.E., 2014, A deposit model for carbonatite and peralkaline intrusion-related rare earth element deposits: U.S. Geological Survey Scientific Investigations Report 2010-5070-J, 58 pp.
- Woolley, A. R. and Kempe, D.R.C., 1989, Carbonatites: Nomenclature, average chemical compositions and element distribution; in Bell, K., ed., *Carbonatites: Genesis and Evolution*: Unwin Hyman, London, pp. 1-14.
- Zech, R. S., Reynolds, R. L., Rosenbaum, J. G., and Brownfield, I. K., 1994, Heavy-mineral placer deposits of the Ute Mountain Ute Indian Reservation, southwestern Colorado and northwestern New Mexico: U.S. Geological Survey, Bulletin 2061-B, 39 pp.

Gallery of Geology

Bastnäesite



Bastnäesite [chemical formula is $(\text{Ce},\text{La})(\text{CO}_3)\text{F}$] in the Red Cloud deposit, Gallinas Mountains district, Lincoln County, New Mexico (length is ~8 mm). Bastnäesite is the most common REE mineral mined in the world today.