



**SME** Society for  
Mining, Metallurgy  
& Exploration

# MINE>XCHANGE



**2024 SME Annual Conference & EXPO**

**FEBRUARY 25-28, 2024 | PHOENIX, AZ**



# GEOCHEMISTRY OF CRITICAL MINERALS IN MINE WASTES IN NEW MEXICO

Virginia T. McLemore, Evan J. Owen, and Richard Kelley  
*New Mexico Bureau of Geology and Mineral Resources, New Mexico Tech,  
Socorro, NM*



# Acknowledgements

- Funding (past and present)
  - Energy Minerals and Natural Resources Department (Abandoned Mine Lands Bureau), U.S. Department of the Interior, Office of Surface Mining and Reclamation (OSMRE)
  - U.S. Geological Survey (including USGS National Geological and Geophysical Data Preservation Program (NGGDPP) and Earth Mapping Resources Initiative (EARTH MRI) program) (G22AC00510)
  - U.S. Department of Energy, CORE-CM project DE-FE0032051 (2021-2024)
- Numerous M.S. theses
- Professional staff and many students who worked on these projects, especially collecting samples
- Thanks to New Mexico Copper Corp. and Santa Fe Gold Inc. for access

[AML Project \(nmt.edu\)](https://nmt.edu)

# Purpose Of NMBGMR Mine Wastes Program


- Test USGS procedures
- Provide data on districts, mines, and mills in New Mexico
  - Help plan and assess reclamation procedures
  - Determine background concentrations
  - Understand geologic processes
  - Identify and quantify critical mineral potential
  - Compare trace-element concentrations in mined versus undisturbed areas
  - Provide background data that can assist with the planning of future mining operations






# Map of mining districts in New Mexico that have critical minerals


- Coal fields are on next slide
- Other areas in New Mexico, such as high-magnesium dolomites, lithium-bearing playas, and coal fields, are not found in specific mining districts and also have potential for critical minerals (CM)





# Coal fields in San Juan Basin, New Mexico

# Location of areas sampled in the New Mexico mine waste study





# Field Procedures

- Inventory
  - Feature descriptions at each site
- Mapping
  - GPS/sketches/LiDAR
  - Radiometric surveys at Black Hawk
    - Divide rock pile into zones based upon radioactivity
- Water Sampling



Joseph working hard at Jicarilla district!



# Field Procedures-continued

- Waste Rock Sampling
  - Soil samples, rock chip samples
    - composite of cells within each feature
    - select of certain areas
    - profiles where feature eroded
  - Sample microbes
  - Select samples
  - Ore samples
  - Slag samples
- Mineralogy, chemistry





Eroded oxidation profile in a rock pile at Copper Flat



# Results



# Results-continued



# Endowment at Copper Flat Tailings, Hillsboro district




Coordinate System: NAD 1983 UTM Zone 13N  
 Projection: Transverse Mercator  
 Datum: North American 1983  
 False Easting: 600,000.0000  
 False Northing: 0.0000  
 Central Meridian: -106.0000  
 Scale Factor: 0.9996  
 Latitude Of Origin: 0.0000  
 Units: Meter

Endowment (kg)	Bi	Co	Cu	Ga	Te	Zn	Zr
Minimum	960	4,230	256,700	13,350	180	29,900	174,000
Maximum	1,700	7,900	574,200	15,600	360	53,400	210,600
Mean	1,280	6,880	402,900	14,240	300	44,000	194,400

Assumes a bulk density of 1430 kg/m<sup>3</sup>

# Acid Rock Drainage (ARD) plot of waste rock pile at mines examined during the NMBGMR mine wastes project



$$AP(\text{kg CaCO}_3/\text{tonnes})=31.25 \times S(\%)$$

$$NP(\text{total C})=83.3 \times C(\%)$$

$$NNP=NP-AP$$

$$NPR=NP/AP$$

# Important note

- Production of critical minerals from mine wastes will likely depend on if the mine wastes can be reprocessed for the major commodity (Au, Ag, base metals)
  - Economic production of solely critical minerals is probably not possible in most cases
- There will likely still be a remaining waste

# Preliminary conclusions

- A sampling protocol has been developed that can estimate the chemical composition of mine wastes
- Larger mine features are divided into subareas for sampling
- Composite samples are collected from cells within the sample areas and homogenized into a single sample
- Elevated critical minerals are found in some of the mine features in New Mexico and these features should be evaluated for potential re-mining or reprocessing
- The available critical minerals are dependent upon types of deposits
- Possible re-mining of mine wastes could clean up these sites and pay for reclamation



# Questions?



SME  
Society for  
Mining, Metallurgy  
& Exploration

**MINE**XCHANGE  
2024 SME Annual Conference & EXPO