

TEXT

Introduction

The Capitan aquifer of Permian (Guadalupe) age is an important source of the ground water used for municipal, industrial, and agricultural purposes in the vicinity of Carlsbad, N. Mex. In addition, water pumped from this aquifer in southern Lea County, New Mexico, and Winkler and Ward Counties, Texas, is injected into other reservoirs by oil companies to increase the recovery of petroleum from partly depleted oil fields (Hiss, 1971). A knowledge of the dimensions of the Capitan aquifer in southeast New Mexico and west Texas will be useful in programs designed to manage this important resource. The map depicting the position and thickness of the Capitan aquifer will also be helpful to petroleum companies who may wish to control "lost circulation" while penetrating this porous and permeable unit as they explore for oil and gas in underlying strata.

The Capitan aquifer

For all practical purposes, the Capitan aquifer is a lithosome that includes the Capitan and Goat Seep Limestones and most or all of the Carlsbad facies of the Artesia Group as recognized by Meissner (1972) and herein used for the near shelf-edge carbonate facies of the Artesia Group. Some of the shelf-margin carbonate banks or reefs in the upper part of the San Andres Limestone are included within the Capitan aquifer wherever they cannot be readily distinguished from the Goat Seep Limestone and Carlsbad facies.

The position and dimensions of the Capitan aquifer closely agree with the Capitan and Goat Seep Limestones and porous and permeable shelf-margin limestone and dolomite members in the San Andres Limestone (Silver and Todd, 1965, Figs. 12 and 13). However, observations of the geometry and lithologic relationships of the shelf-margin and shelf sediments rocks in the field suggest that the thickness of the Capitan Limestone is considerably less than is usually shown by others who have mapped this stratigraphic reef. The relationships between the now obsolete Carlsbad Limestone and Capitan Limestone expressed by Dunham (1972, Fig. 1-1) appear to more closely match the field relationships observable in the vicinity of Carlsbad and White City, N. Mex.

The Capitan aquifer is generally composed of relatively "clean" carbonate rock, especially near the back-reef edge. The radioactivity recorded on a gamma-ray electrical log of the Capitan and (or) Goat Seep Limestone is characteristically very low as shown in stratigraphic sections A-A' and B-B'.

The tops and bases of the Capitan aquifer were determined primarily on the basis of the vertical extent of the relatively "clean" carbonate rock as indicated by the low gamma-ray activity levels shown on the electrical logs and the general stratigraphic position. Lithologic logs of field scout formations "tops" report of lost circulation, and other information were used whenever available to confirm these determinations. Zones containing 50 percent or less of interbedded back or fore-reef lithofacies were included with the Capitan aquifer as a matter of convenience; therefore, the net aggregate thickness of the Capitan aquifer may have been increased slightly by this arbitrary procedure.

It is often difficult or impossible to distinguish between other reefs and carbonate mounds in the back-reef sediments and the Capitan and Goat Seep Limestones solely on the basis of the responses recorded on gamma-ray, sonic, and neutral electrical logs. Adjacent to the shelf edge, the Capitan and Goat Seep Limestones in the Carlsbad facies of the Artesia Group are included as part of this aquifer wherever the chemical composition of water, the water-level declines, or the natural radioactivity in the adjacent units are similar to that in the main aquifer (Hiss, 1973 and 1974a).

Units previously described as reefs of Yates or Seven Rivers age, part of the Grayburg Formation, and the shelf-margin carbonate banks in the upper part of the San Andres Limestone are considered to be part of the Capitan aquifer wherever they cannot be distinguished as separate entities and the water quality, electrical log characteristics, or hydraulic response justify inclusion.

Dimensions of the Capitan aquifer

Determination of extent and thickness

The locations of nearly 400 wells that have been drilled through the Capitan aquifer within the project area, along with the corresponding thickness of the Capitan aquifer penetrated in each well, are plotted on the map. Gamma-ray, gamma-ray-neutron, or other combinations of electrical logs of the Capitan aquifer were used to determine the thickness of these wells. Lithologic logs were available for approximately 15 percent of the wells.

Lateral extent

The Capitan aquifer parallels the north and east margins of the Delaware basin in an arcuate strip extending from the Guadalupe Mountains southwest of Carlsbad, N. Mex. to the Glass Mountains southwest of Fort Stockton, Tex. Outcrops of the Capitan Limestone have been found in the Glass, Guadalupe, Apache, and Delaware Mountains. Undoubtedly it is present elsewhere in the subsurface along the west and southwest margins of the Delaware basin.

As shown on the map, the Capitan aquifer is one continuous unit along the north and east margins of the Delaware basin and the Guadalupe Mountains. The Capitan aquifer by faulting appears to be limited to the mountainous areas along the western and southern parts of the Delaware basin. Faults have not been observed in the subsurface along the western edge of the Central Basin platform and the southern edge of the Northwestern shelf. The irregular top and bottom surfaces and the lobate and back-reef edges are depositional forms (map and stratigraphic sections A-A' and B-B') that might appear as a major lateral displacement of the Capitan aquifer in the vicinity of T-22 S., R-25 E., approximately 12 miles (24 km) southwest of Carlsbad, is not a post-Capitan age fault. This shift in position is probably due to growth of the Capitan reef. A pre-Guadalupe age fault-controlled alignment of the Delaware basin (Hiss, 1963, p. 1715; and 1970).

The width of the Capitan aquifer ranges from about 10 to more than 14 miles (16 to 23 km) along the edge of the Northwestern shelf in the vicinity of Carlsbad to the central part of southern Lea County, New Mexico. The Capitan aquifer is much more restricted in the western edge of the Central Basin platform, where it seldom exceeds 11 miles (18 km) in width. The fore-reef edge of the Capitan aquifer appears to be relatively steep throughout the area. If exposed, this abrupt slope would undoubtedly resemble the reef escarpment southwest of Carlsbad in the Delaware Mountains (Hiss and others, 1964; Newell and others, 1953). Well control is adequate for definition of the fore-reef edge of the Capitan aquifer at several locations in the subsurface. Approximately 1,200 feet (366 m) of vertical relief along the fore-reef edge of the Capitan aquifer was detected on two oil wells drilled within a few hundred feet of horizontal distance in secs. 5 and 9, T-22 S., R-23 E., Lea County, New Mexico (Meissner, 1972, plate 12). Similar evidence of the relative steepness of the fore-reef side of the Capitan aquifer can be obtained where deep-well drilling is concentrated in the newly discovered gas fields in the vicinity of Pecos, Wink, and Fort Stockton in Winkler, Ward, and Pecos Counties, Texas, respectively.

The back-reef edge of the Capitan aquifer is much more irregular than the fore-reef edge and is gradual. In some areas, especially along the western edge of the northern part of the Central Basin platform, it is difficult to distinguish the Capitan aquifer from the upper part of the San Andres Limestone. In this area the Capitan aquifer has been extended to include the stratigraphic reef facies of the San Andres Limestone because of the proximity, similar lithology, and hydraulic behavior of the two units (stratigraphic section B-B').

Thickness

As is readily observed from the map, the thickness of the Capitan aquifer is quite variable. The Capitan aquifer appears to be composed of irregularly shaped and spaced thick accumulations of carbonate rock alternating with thinner units composed of similar material. Many of the locally thick areas are behind the reef front and are composed of carbonate banks, islands, or mounds which flourished behind the protection of the reef crest (Kendall, 1965, p. 259 and pls. 2 and 3). Hiss (1968 and 1972) has mapped and described both current-oriented and irregularly oriented "shelf down" carbonate mounds in the vicinity of Dark Canyon southwest of Carlsbad, N. Mex.

A number of small oil fields located along the trend of the Capitan aquifer are apparently localized on carbonate "buildups" that have been referred to by Stipp and Haigler (1956) as "reef knobs" interspersed between "large channels." The majority of these carbonate mounds or "buildups" are also located within the thick areas shown in the Capitan aquifer thickness map. The Capitan aquifer attains a maximum thickness of 2,357 feet (718 m) in the Odessa Natural Gas Federal Dooly well located on one of these mounds in sec. 24, T-20 S., R-29 E., about 13 miles (21 km) northeast of Carlsbad.

Hiss (1974b) has identified several of the most prominent linear, thin areas oriented transverse to the arcuate main trend of the Capitan aquifer as submarine canyons. The submarine canyons are contemporary in age to the reefs that comprise the Capitan aquifer.

The Capitan aquifer is slightly thicker along the edge of the Northwestern shelf in New Mexico than in Texas. In addition, the areal extent of the thick areas is correspondingly larger. These differences probably reflect the shape of the structural framework of the Delaware basin at the time of the deposition of the Capitan aquifer. The more extensive reefs were developed along the margin of a broad, shallow shelf on the northern part of the basin, whereas narrow, linear reefs grew along the faulted western margin of the Central Basin platform.

Acknowledgments

This map was prepared in cooperation with the New Mexico State Engineer. Many of the data were generously provided by several oil companies, consultants, and private individuals. Mr. George J. Galt assisted the author with the preparation of the map and stratigraphic sections.

References

Dunham, R. J., 1972, Capitan reef, New Mexico and Texas: Facts and questions to aid interpretation and group discussion: Soc. Econ. Paleontologists and Mineralogists, Permian Basin, Sec. 1, Pub. 72-14, approx. 200 p.

Green, W. R., and others, eds., 1964, Geology of the Capitan reef complex of the Guadalupe Mountains, Culberson County, Texas and Eddy County, New Mexico: Roswell Geol. Soc. Guidebook, Field trip, May 6-9, 1964, 124 p.

Hiss, J. M., 1963, Late Paleozoic tectonics and mountain ranges, western Texas to southern Colorado: Am. Assoc. Petroleum Geologists Bull., v. 47, no. 9, 1709-1725, 6 figs.

—1970, Late Paleozoic structural directions in southern Permian basin, west Texas and southeastern New Mexico: Am. Assoc. Petroleum Geologists Bull., v. 54, no. 10, p. 1809-1827, 16 figs.

Hiss, W. L., 1971, Water-level changes in the Permian Capitan aquifer, Eddy and southern Lea Counties, New Mexico: New Mexico Prof. Engineer, November, 1971, p. 16-22.

—1973, Capitan aquifer observation-well network, Carlsbad to Jal, New Mexico: New Mexico State Engineer Tech. Rept. no. 38, 76 p., 11 figs.

—1974a, Map showing chloride-ion concentration in water from sedimentary rocks of Permian (Guadalupe) age, southeastern New Mexico and western Texas: U.S. Geol. Survey open-file report.

—1974b, Reduction of ground-water flow through the Capitan and Goat Seep Limestones by incised contemporaneous submarine canyons [abs.]: G. C. M. B. A. Survey, In Abstracts with programs, October 1974, v. 6, no. 7, p. 797.

Kendall, C. G. St. C., 1969, An environmental re-interpretation of the Permian evaporite-carbonate shelf sediments of the Guadalupe Mountains: Geol. Soc. America Bull., v. 80, no. 12, p. 2503-2536.

Meissner, F. F., 1972, Cyclical sedimentation in mid-Permian strata, in Elam, J. G., and Chubb, Stewart, eds., Cyclical sedimentation in the Permian basin, and other: West Texas Geol. Soc. pub. 72-60, p. 203-232.

Notts, W. S., 1968, The control of ground-water occurrence by lithofacies in the Guadalupe reef complex near Carlsbad, New Mexico: Geol. Soc. America Bull., v. 79, p. 283-298.


—1972, Geology and paleoenvironments of the northern segment, Capitan shelf, New Mexico and west Texas: Geol. Soc. America Bull., v. 83, no. 3, p. 701-722.

Newell, N. D., and others, 1953, The Permian reef complex of the Guadalupe Mountains region, Texas and New Mexico—a study in paleogeology: San Francisco, W. H. Freeman & Co., 236 p.


Oriel, S. S., Myers, D. A., and Crosby, E. J., 1967, West Texas Permian basin region, in McKee, E. D., and Oriel, S. S., eds., Paleontological Investigations of the Permian System in the United States: U.S. Geol. Survey Prof. Paper 519, p. 17-62.

Silver, B. A., and Todd, R. G., 1969, Permian cyclic strata, northern Midland and Delaware basins, west Texas and southeastern New Mexico: Am. Assoc. Petroleum Geologists Bull., v. 53, no. 11, p. 2223-2251.


Stipp, T. F., and Haigler, L. B., 1956, Preliminary structure contour map of a part of southeastern New Mexico showing oil and gas development: U.S. Geol. Survey Oil and Gas Inv. Map 1077.




NEW MEXICO base from U.S. Geological Survey, Branch of Oil and Gas Operations, Base 11.9 South of 2 miles (1961). Texas base from Midland Map Co. #2 (1961), #3 (1961), #4 (1961), #5 (1961), #6 (1955), #7 (1961), #8 (1961), #9 (1955), and #10 (1955) 1 inch = 5,000 feet, used with permission of Midland Map Co.



TRANSVERSE STRATIGRAPHIC SECTION A-A' SHOWING THE POSITION OF THE CAPITAN AQUIFER IN THE CENTRAL PART OF SOUTHERN LEA COUNTY, NEW MEXICO



TRANSVERSE STRATIGRAPHIC SECTION B-B' SHOWING THE POSITION OF THE CAPITAN AQUIFER IN NORTHERN WARD COUNTY, TEXAS



WELL IDENTIFICATION
 ORDER OF ELEMENTS:
 OPERATOR, lease, well number,
 geographic location, reference
 datum, total depth.

THICKNESS OF THE PERMIAN GUADALUPE CAPITAN AQUIFER, SOUTHEAST NEW MEXICO AND WEST TEXAS
 by W. L. HISS