

Geologic Map of the Santa Fe 7.5-minute quadrangle

by Adam S. Read, Daniel J. Koning, Gary A. Smith, Stever Raiser, John Rogers, and Paul W. Bauer
May, 2000
Last Revised: 8-October-2003
1:24,000

COMMENTS TO MAP USERS
A geologic map displays information on the distribution, nature, orientation, and age relationships of rock deposits and the occurrence of structural features. Geologic and fault contacts are shown on the map. Boundaries between different types of units. Data depicted on this geologic quadrangle map may be based on any of the following: reconnaissance field geologic mapping, compilation of published and unpublished work, and photogeologic interpretation. Locations of contacts are not surveyed, but are plotted by interpretation of the position of a given contact on a topographic base map; therefore, the accuracy of locations depends on the scale of mapping and the interpretation of the geologists. Any enlargement of this map could cause misunderstanding in the detail of mapping and may result in erroneous interpretations. Site-specific conditions should be verified by detailed surface mapping or subsurface exploration. Topographic and cultural changes associated with recent development may not be shown.

This work was performed under the STATEMAP component of the USGS National Cooperative Geologic Mapping Program. Funding was provided by the U.S. Geological Survey and the New Mexico Bureau of Geology and Mineral Resources, a division of New Mexico Tech.

Description of Map Units

Grain sizes follow the Lithic-Wentworth scale for clastic sediments (Sklar, 1944; Wentworth, 1922) and are based on field estimates. Sand is classified according to Penland et al. (1987). The term "clastic" refers to the size factor (particles less than 2 mm in diameter). Percentages are based on percent volume and were estimated in the field with the aid of percentage charts. Colors of nonconsolidated sediments are based on visual comparison of dry samples to the Munsell Soil Color Charts (Munsell Color, 1953). Surface slickensides are color-coded on the map if they are estimated to be at least 1 m thick. Soil horizon designations and descriptive terms follow those of the Soil Survey Staff (1993) and Birkeland et al. (1991). Stages of pedogenic calcium carbonate morphology follow those of Gile et al. (1966).

Cenozoic Rocks
Quaternary Hillside Deposits
Qes Collium and sheetwash deposits (Middle?) Pleistocene to Holocene - Generally light yellowish brown to yellowish brown (10YR 5-6.5) gravel, sand, silt, and minor clay. Gravel is angular to subangular, poorly to moderately sorted, finely matrix-supported and generally cobbles and pebbles. Bedding is non-existent or very fine, thin, and lenticular. Sediment is loose to weakly consolidated and 1 to 12.7 m thick. Unit includes minor alluvium.

Quaternary Anthropogenic Deposits
Qaf Artificial fill (Historical) - Brown (10YR 5/3) silt, sand, and gravel. Sediment is poorly sorted and lacks bedding. Sand is subdivided to subangular and an arkosic arenite. Sediment is loose and of variable thickness. Unit includes minor alluvium.

Quaternary Alluvial Deposits
Qol (labeled Qo also) Alluvium in modern drainage channels (Active) - Sand, pebbles, cobbles, and boulders from an active drainage channel (Qol) at 58-31.1 or younger (Qol). Sediment generally consists of light yellowish brown to light brown (7.5YR 10/4-6) sandy gravel. Clay is supported and is very fine to very coarse-grained, moderately sorted, and angular to subangular cobbles and pebbles. It is poorly sorted and is in the mountain. In the Big Tesaque watershed there is at least one of these units was mapped where there is a significant increase in slope.

Qom Younger alluvial fan deposits (uppermost Pleistocene to Holocene) - Alluvial alluvium at the mouth of small tributary drainage in the Santa Fe River. The upper portion of these fans grade out over Qol at 58-31.1 or younger (Qol). Sediment generally consists of light yellowish brown to light brown (7.5YR 10/4-6) sandy gravel and generally sand plus minor silt. Sediment is clay-supported and is very fine to very coarse-grained, moderately sorted, and angular to subangular cobbles and pebbles. It is poorly sorted and is in the mountain. In the Big Tesaque watershed there is at least one of these units was mapped where there is a significant increase in slope.

Qon Terrace deposits above Arroyo Honda (Pleistocene to Holocene)
Qn1 Terrace deposit (middle Pleistocene) - Fill terrace gravel with reads approximately 50 to 60 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn2 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 35 to 40 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn3 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 20 to 25 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn4 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 15 to 20 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn5 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 10 to 15 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn6 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 5 to 10 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn7 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn8 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn9 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.



Qn10 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn11 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn12 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn13 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.

Qn14 Terrace deposit (middle Pleistocene) - Fill terrace gravel with a read approximately 0 to 5 m above modern grade along the Big Tesaque. In many places it is clear that the base of this gravel deposit is below the elevation of the modern valley floor. The maximum thickness of the fill is unknown. Base exposures along steep rises reveal a complicated fault stratigraphy of alternating cobble-to-boulder gravel and pebbly sand with rare interstratified thin beds of fine-grained siltstone and claystone.



Comments on Faults in the Vicinity of Bishop's Lodge Arroyo
Number Member (Upper Pleistocene?) to lower Miocene - Poorly sorted sand to cobble conglomerates, sandstone, and minor mudstone composed of detrital rocks from pre-Tertiary rocks. Color is typically red to pink but is locally white to very pale pink and white.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

References Cited
Buchanan, C.O., and Nelson, H.H., 1978. New K-Ar dates and the late Pleistocene to Holocene geomorphic history of the central Rio Grande region, New Mexico. Geological Society of America Bulletin, vol. 89, p. 231-247.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.


General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.



Comments on Faults in the Vicinity of Bishop's Lodge Arroyo
Number Member (Upper Pleistocene?) to lower Miocene - Poorly sorted sand to cobble conglomerates, sandstone, and minor mudstone composed of detrital rocks from pre-Tertiary rocks. Color is typically red to pink but is locally white to very pale pink and white.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

General Observations on Faults in the Arroyo Formation: Faults are generally not obvious within the Number Member unless anomalous bedding attitudes are present and recognizable. Where obvious, they are generally normal faults, such as those of the Arroyo and the central Rio Grande rift. They are generally not as well developed as those of the central Rio Grande rift.

