

Introduction

This geologic compilation map was produced as part of a regional hydrogeologic study of the Southern Sacramento Mountains. New geologic mapping was limited to the area east of the crest of the Sacramento Mountains, where the San Andres and Yano Formations are exposed. Work was begun in late 2005 and completed in early 2009. Comparison of map data sets and cartographic processing were completed in early 2012. The new mapping was completed together with the mapping of Pray (1961) of the western escarpment of the range and some northern portions of the map block (1973), as shown on the data source index map.

Methodology

In addition to standard geologic field mapping procedures, much geology, especially in the sparsely wooded eastern portion of the study area, was interpreted from aerial photographs. Substitution of the members of the San Andres formation into its constituent members (as defined by Kelley, 1971) was performed in this way. These interpretations were locally field checked and compared to the regional map of Kelley (1971). In most places, the interpretations are similar. Pray (1961) mapped the Yano Formation to include the 'Hondo Member' of the San Andres formation, but in the study area the Yano Formation overlies basal limestone beds of the San Andres up to and including the distinctive tan quartz sandstones that are equivalent to the Gloria Sandstone of northern New Mexico. This member ranges from 80 to 120 feet thick (Pray, 1961) and is generally considered the only reliably mappable horizon above the Abso-Yano contact. The Yano contact directly beneath the tan sandstone beds, and slightly thicker and more gradational in appearance than the underlying Bonney Canyon member. Although it contains abundant gypsum elsewhere (Kelley, 1971), in the mapped area it is composed of thin bedded, locally brecciated dolomite (Ziegler, personal communication, 2007). The total thickness is not exposed, but it is at least 100 feet.

Yano and San Andres Formations

Geology on the map west of a roughly north-south through Rio Hondo (approximately following the West Side Road), is largely from Pray (1961). Aside from basic unit descriptions provided here, the reader is referred to his excellent discussion of the geology and the stratigraphy and structures of this area. From the aforementioned line to the east of the range and east the exposed rocks are the Permian Yano and San Andres Formations, which dip shallowly (2-3°) to the east. The new mapping presented here is largely concerned with these units, so additional discussion is warranted. Good natural exposures of the Yano Formation are rare, as it is usually covered with colluvium from the overlying, more resistant San Andres Formation or by valley bottom alluvium. A complete section is not exposed east of the range crest. Pray (1961) measured a complete section 1230 feet thick on the west flank of Crowder Peak three miles west of Timmeron at the southeast corner of the present study area. Estimated on this section and the Southern Production Company #1 oil well west between Cloudcroft and the Rio Pecos, Pray (1961) estimated the Yano Formation at 1200 to 1800 feet in the western Sacramento Mountains. The differences are probably due to local stratigraphic variation and erosion of fragments. Good natural exposures of the Yano Formation are rare, as it is usually covered with colluvium from the overlying, more resistant San Andres Formation or by valley bottom alluvium. A complete section is not exposed east of the range crest. Pray (1961) measured a complete section 1230 feet thick on the west flank of Crowder Peak three miles west of Timmeron at the southeast corner of the present study area. Estimated on this section and the Southern Production Company #1 oil well west between Cloudcroft and the Rio Pecos, Pray (1961) estimated the Yano Formation at 1200 to 1800 feet in the western Sacramento Mountains. The differences are probably due to local stratigraphic variation and erosion of fragments. Good natural exposures of the Yano Formation are rare, as it is usually covered with colluvium from the overlying, more resistant San Andres Formation or by valley bottom alluvium. A complete section is not exposed east of the range crest. Pray (1961) measured a complete section 1230 feet thick on the west flank of Crowder Peak three miles west of Timmeron at the southeast corner of the present study area. Estimated on this section and the Southern Production Company #1 oil well west between Cloudcroft and the Rio Pecos, Pray (1961) estimated the Yano Formation at 1200 to 1800 feet in the western Sacramento Mountains. The differences are probably due to local stratigraphic variation and erosion of fragments.

CORRELATION OF UNITS

Historic	Quaternary	Tertiary	Permian	Pennsylvanian	Mississippian	Silurian/Devonian	Ordovician	Precambrian
daf	Qa	Qs	Qg	Tl	Psf	Psb	Pr	Py
Qa	Qs	Qg	Tl	Psf	Psb	Pr	Py	Py
Qs	Qg	Tl	Psf	Psb	Pr	Py	Py	Py
Qg	Tl	Psf	Psb	Pr	Py	Py	Py	Py
Tl	Psf	Psb	Pr	Py	Py	Py	Py	Py
Psf	Psb	Pr	Py	Py	Py	Py	Py	Py
Psb	Pr	Py	Py	Py	Py	Py	Py	Py
Pr	Py	Py	Py	Py	Py	Py	Py	Py
Py	Py	Py	Py	Py	Py	Py	Py	Py
Pa	Pb	Pg	Mu	Sdu	Ou	pCu	Py	Py
Pb	Pg	Mu	Sdu	Ou	pCu	Py	Py	Py
Pg	Mu	Sdu	Ou	pCu	Py	Py	Py	Py
Mu	Sdu	Ou	pCu	Py	Py	Py	Py	Py
Sdu	Ou	pCu	Py	Py	Py	Py	Py	Py
Ou	pCu	Py	Py	Py	Py	Py	Py	Py
pCu	Py	Py	Py	Py	Py	Py	Py	Py
Py	Py	Py	Py	Py	Py	Py	Py	Py

DESCRIPTION OF UNITS



- Qa** Undivided valley stream alluvium, valley fill, and aeolian deposits (Quaternary)—Stream alluvium consists of unsorted gravel and poorly to moderately sorted clay, silt, sand in active stream channels and ephemeral arroyos. Valley fill is composed of poorly to moderately sorted clay, silt, and sand, often distinctly stratified and commonly with angular to subangular clasts of bedrock. Matrix material is light to dark brown. Valley fill grades into alluvial and colluvial fans on toes of alluvial fans.
- Qs** Slump blocks, landslide deposits, and associated colluvium (Quaternary and Tertiary)—Homoclinal slope failure deposits and rotational landslide blocks in the central portion of the map area. Stabilized by vegetation and not active.
- Qg** Terrace gravels, alluvial fans, and upland deposits, undivided (Quaternary and Tertiary)—Generally coarse deposits composed largely of sand and gravel and incised by modern drainages. Extensive gravel terraces in the eastern and southern portions of map area, and lower terraces of above modern drainages, and may be as old as Miocene. Thickness: 0-60 ft.
- Tl** Intra-Permian igneous rocks (Tertiary)—Sills and subordinate dikes of porphyritic andesite to latite. Thickness: dikes a few feet to tens of feet; sills up to 200 ft.
- Psf** Four-mile Draw member, San Andres Formation (Permian)—Thin bedded, locally brecciated grayish-brown dolomite. Distinctly darker on aerial photographs than underlying members. Contains abundant gypsum outside of the study area. Thickness: top of unit exposed, 1-100 ft.
- Psb** Bonney Canyon member, San Andres Formation (Permian)—Light to dark gray, bluish-gray, and tan thin- to medium-bedded dolomite and limestone. Thin bedded. This bedding forms distinctive 'brix-eye' pattern on aerial photos. Thickness: 200 - 400 ft.
- Pr** Rio Bonito Member, San Andres Formation (Permian)—Light to dark gray, bluish-gray, and tan medium- to thick-bedded dolomite and limestone. Distinctive Hondo Sandstone member 40 to 80 feet above base of unit is composed of brown to tan clean quartz sandstone with well-sorted, rounded grains and is equivalent to the Gloria Sandstone of central and northern New Mexico. Thickness: 400-600 ft.
- Py** Yano Formation (Permian)—Yellow to tan siltstone and fine sandstone, red to pink, mudstone, and fine sandstone, and gray to tan, often silty, dolomite and subordinate limestone. Gypsum, anhydrite, and halite are present below 1000 ft in deep test wells but have been removed by dissolution near to the surface, resulting in abundant collapse features and chaotic bedding. Thickness: highly variable, ~1500 ft in deep well near Cloudcroft, thin to 0 at Pagito Mountain on the Mesalero Reservation.
- Pa** Abso Formation (Permian)—Dark reddish-brown mudstone, anoxic sandstone, and local basal conglomerate. Southeastward thinning of thin-bedded limestone, dolomite and gray shale in southern portion of map area. Thickness: 200-500 ft.
- Pb** Barum Formation (Pennsylvanian)—Red and gray sandstone and shale, gray limestone, and limestone conglomerate. Red beds and conglomerate become more abundant to the south. Thickness: Three forms 350 to 0 ft north to south, and an angular unconformity between Pennsylvanian strata and the overlying Abso Formation.
- Pg** Helder Formation (Pennsylvanian)—Cyclic gray limestones, gray and red calcareous shale, mudstone, sandstone and conglomerate. Massive, cliff-forming bioherms and biotremal limestones present near base of unit. Thickness: 0-850 ft.
- Mu** Goodler Formation (Pennsylvanian)—Basal coarse quartzite sandstone and dark argillaceous silty, and cherty limestone. These units are overlain by thick bedded to massive cherty limestone of the Bug-Scuffs limestone member, which grades and interfingering laterally to with equivalent detrital facies composed of interbedded sandstone, siltstone, shale and minor limestone. Thickness: 1200-1800 ft.
- Sdu** Callalero, Lake Valley, Rancharia, and Helms Formations, undivided (Mississippian)—Gray and dark gray, locally cherty or silty limestone, marl and argillaceous and calcareous shale. Bioherms up to 30 ft thick are prominent in the lower Lake Valley Formation. Thickness: 350-450 ft.
- Ou** Fossiliferous, Otero, Sly Gap and Pecha (?), Formations, undivided (Silurian and Devonian)—Dark gray, olive and brownish-gray cherty dolomite, medium gray to brownish gray and olive silty dolomite, dolomitic quartz sandstone and sandstone, gray calcareous shale, mottled gray limestone, and dark gray to light gray shale. Thickness: 135-175 ft.
- pCu** Precambrian rocks, undivided—Gray-green shale, siltstone, and minor quartz sandstone intruded by diabase. Thickness: at least 80 ft exposed.
- Py** Precambrian rocks, undivided—Gray-green shale, siltstone, and minor quartz sandstone intruded by diabase. Thickness: at least 80 ft exposed.

COMMENTS TO MAP USERS **24K QUAD INDEX MAP AND GEOLOGIC MAPPING CREDITS**

A geologic map displays information on the distribution, nature, orientation, and age relationships of rock and deposits and the occurrence of structural features. Geologic and fault contacts are irregular surfaces that form boundaries between different types of geologic units. Data reported on this geologic quadrangle map may be based on any of the following reconnaissance field geologic mapping, compilation of published and unpublished work, and photogeologic interpretation. Locations of contacts are not surveyed, but are plotted by interpretation of the position of a given contact onto a topographic base map; therefore, the accuracy of contact locations depends on the scale of mapping and the interpretation of the geologist. An enlargement of this map could cause misunderstanding in the detail of mapping and may result in erroneous interpretations. Site-specific conditions should be verified by detailed surface mapping or subsurface exploration. Topographic and cultural changes associated with recent development may not be shown.


Cross sections are constructed based upon the interpretations of the author made from geologic mapping, and available geophysical, and subsurface (drillhole) data. Cross-sections should be used as an aid to understanding the general geologic framework of the map area, and not be the sole source of information for use in locating or designing wells, buildings, roads, or other man-made structures.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the State of New Mexico, or the U.S. Government.



MAP EXPLANATION

Geologic contact	Fault, exposed	Anticline, exposed	Horizontal bedding
Cross section line	Fault, intermittent-obscured	Anticline, concealed	Inclined bedding
	Fault, concealed	Syncline, exposed	
	Fault on downthrown side	Syncline, concealed	
	Strike slip fault, dextral	Monoclinal, exposed	
		Monoclinal, concealed	



New Mexico Bureau of Geology and Mineral Resources
Open-File Report 537-Geologic Map, Sheet 1

Generalized Geologic Map of the Southern Sacramento Mountains, Otero and Chaves Counties, New Mexico

February, 2012
by **Geoffrey Rawling**

New Mexico Bureau of Geology and Mineral Resources
801 Leroy Place, Socorro, New Mexico 87801-4796
[575] 835-5490
http://geoinf.nm.edu

COOPERATIVE GEOLOGIC MAPPING PROGRAM