

K-Ar age of alunite alteration at Red Mountain, Lake City area, western San Juan Mountains, Colorado

H.H. Mehnert, J.F. Slack, and G.T. Cebula

Isochron/West, Bulletin of Isotopic Geochronology, v. 28, pp. 9

Downloaded from: <https://geoinfo.nmt.edu/publications/periodicals/isochronwest/home.cfm?Issue=28>

Isochron/West was published at irregular intervals from 1971 to 1996. The journal was patterned after the journal *Radiocarbon* and covered isotopic age-dating (except carbon-14) on rocks and minerals from the Western Hemisphere. Initially, the geographic scope of papers was restricted to the western half of the United States, but was later expanded. The journal was sponsored and staffed by the New Mexico Bureau of Mines (now Geology) & Mineral Resources and the Nevada Bureau of Mines & Geology.



ISOCHRON/WEST
A Bulletin of Isotopic Geochronology

All back-issue papers are available for free: <https://geoinfo.nmt.edu/publications/periodicals/isochronwest>

This page is intentionally left blank to maintain order of facing pages.

K-AR AGE OF ALUNITE ALTERATION AT RED MOUNTAIN, LAKE CITY AREA, WESTERN SAN JUAN MOUNTAINS, COLORADO

HARALD H. MEHNERT
JOHN F. SLACK
GERALD T. CEBULA

U.S. Geological Survey, Denver, CO 80225
U.S. Geological Survey, Reston, VA 22092
U.S. Geological Survey, Denver, CO 80225

The San Juan Mountains of southwestern Colorado are famous for rich vein deposits of base and precious metals and for patterns of conspicuous hydrothermal alteration. Areas of acid-sulfate alteration, characterized by extensive base leaching of volcanic rocks or hypabyssal plutons, are of current interest because of their local association with Au, Ag, and, in some places, U mineralization (Lipman and others, 1976; Cunningham and others, 1978; Steven and others, 1979). Potential may also exist for recovery of aluminum from altered rocks rich in alunite (Hall, 1978). Alunite has been shown to be suitable for K-Ar dating in other mining districts (Mehnert and others, 1973b; Ashley and Silberman, 1976). We report here two concordant K-Ar ages determined on hydrothermal alunite from Red Mountain, in the Lake City area of the western San Juan Mountains.

GEOLOGIC SETTING

The San Juan Mountains comprise a large continental volcanic field chiefly of middle Tertiary age (Larsen and Cross, 1956; Lipman and others, 1970). Initial volcanism, in early Oligocene time, was dominated by andesitic strato-volcanoes that produced a coalesced platform of lavas, breccias, and mudflows. During the late Oligocene and Miocene, silicic ash-flow tuffs were erupted from at least 15 caldera sources (Steven and Lipman, 1976). The youngest caldera is Lake City caldera, in the western San Juan Mountains (Lipman and others, 1973). The Lake City caldera formed 22.5 m.y. (million years) ago, in response to pyroclastic eruptions of the Sunshine Peak Tuff; the age assignment is based on K-Ar dating of both the intracaldera and outflow tuff members (Mehnert and others, 1973a).

Red Mountain is a topographically prominent flow dome that was emplaced along the eastern ring-fault zone of the Lake City caldera (fig. 1). Extensive talus and poor exposures prevent detailed mapping, but the dome is believed to extend downward into a volcanic neck (Lipman, 1976). Most of the dome appears to be variably altered quartz latite porphyry, containing phenocrysts of quartz, sanidine, and plagioclase. Hydrothermal alteration is locally pervasive and has in some places destroyed primary volcanic textures. Major alteration minerals are quartz, alunite, and kaolinite. The alunite occurs as irregular disseminations, as replacements of feldspar phenocrysts, or, less commonly, as thin veinlets. Alunite for isotopic dating was obtained from drill core furnished by Earth Sciences, Inc., Golden, Colorado (see fig. 1 for drill-hole location). K-Ar ages were

determined on a sample of massive alunite from a cross-cutting veinlet in the core at a depth of 647 feet, and on a core sample of disseminated alunite from 683 feet. The K-Ar data yield concordant ages of 23.3 m.y. \pm 1.1 m.y. and 22.9 m.y. \pm 1.6 m.y., respectively.

DISCUSSION

The K-Ar dates on alunite from Red Mountain indicate that the hydrothermal alteration there was coeval with the Lake City caldera cycle. They also agree, within analytical uncertainty, with a mean K-Ar age of 22.8 m.y. \pm 0.6 m.y. determined for an unaltered lava flow that overlies altered flows probably related to venting of the Red Mountain dome (Mehnert and others, 1973a). The acid-sulfate alteration assemblage at Red Mountain is considered to be a shallow facies of the same hydrothermal system that deposited base and precious metals, as well as local U (at the Golden Fleece mine, fig. 1), in nearby fissure veins (Slack and Lipman, 1979). The entire system developed during the waning stages of the caldera cycle, and the hydrothermal solutions probably vented as solfataric hot springs above the present crest of Red Mountain.

SAMPLE DESCRIPTIONS

Fresh alunite for isotopic dating was obtained from drill core furnished by Earth Sciences, Inc., Golden, Colo. Mineral separations and analyses were carried out in the laboratories of the U.S. Geological Survey in Denver, Colo. Constants used in calculating the K-Ar age are:

$${}^{40}\text{K}\lambda_{\epsilon} = 0.581 \times 10^{-10}/\text{y}, \quad {}^{40}\text{K}\lambda_{\beta} = 4.962 \times 10^{-10}/\text{y}, \quad {}^{40}\text{K}/\text{K} = 1.167 \times 10^{-4}. \quad \text{Analytical uncertainty is quoted as } 2\sigma.$$

1. *U.S.G.S.(D) -DKA3633* K-Ar
Quartz latite from Red Mountain dome, Lake City caldera (Hinsdale County, CO; 37°58'N, 107°20'W). Disseminated alunite replacement of feldspar phenocrysts in drill core from depth of 683 feet. *Analytical data:* K₂O = 4.54%, 4.55%; ⁴⁰Ar = 1.508 \times 10⁻¹⁰ moles/gram; ⁴⁰Ar/Ar = 38.5%. *Analyzed by:* H. H. Mehnert. *Comment:* Time of hydrothermal alteration. (alunite) 22.9 m.y. \pm 1.6 m.y.




FIGURE 1. Sketch map of the Lake City caldera and vicinity, showing veins and principal ore assemblages (from Slack and Lipman, 1979). Drill hole is near the summit of Red Mountain, in the center of the mapped area of quartz latite porphyry.

2. *U.S.G.S.(D) -DKA3635* K-Ar
 Quartz latite from Red Mountain dome, Lake City
 caldera (Hinsdale County, CO; 37°58'N, 107°20'W).
 Fine-grained vein alunite from drill core at depth of 647
 feet. *Analytical data*: K₂O = 8.76%, 8.79%; ⁴⁰Ar =
 2.956 × 10⁻¹⁰ moles/gram; ⁴⁰Ar/Ar = 60.7%. *Ana-*
lyzed by: H. H. Mehnert. *Comment*: Time of hydro-
 thermal alteration.

(alunite) 23.3 m.y. ± 1.1 m.y.

REFERENCES

- Ashley, R. P., and Silberman, M. L. (1976) Direct dating of mineralization at Goldfield, Nevada, by potassium-argon and fission-track methods: *Econ. Geol.*, v. 71, p. 904-924.
- Cunningham, C. G., Steven, T. A., and Naeser, C. W. (1978) Preliminary structural and mineralogical analysis of the Deer Trail Mountain-Alunite Ridge mining area, Utah: U.S. Geol. Survey open-file rept. 78-314.
- Hall, R. B. (1978) World nonbauxite aluminum resources—alunite: U.S. Geol. Survey Prof. Paper 1076-A.
- Larsen, E. S., Jr., and Cross, W. (1956) Geology and petrology of the San Juan region, southwestern Colorado: U.S. Geol. Survey Prof. Paper 258.
- Lipman, P. W. (1976) Geologic map of the Lake City caldera area, western San Juan Mountains, southwestern Colorado: U.S. Geol. Survey Misc. Inv. Map I-962, scale 1:48,000.
- Lipman, P. W., Fisher, F. S., Mehnert, H. H., Naeser, C. W., Luedke, R. G., and Steven, T. A. (1976) Multiple ages of mid-Tertiary mineralization and alteration in the western San Juan Mountains, Colorado: *Econ. Geol.*, v. 71, p. 571-588.
- Lipman, P. W., Steven, T. A., Luedke, R. G., and Burbank, W. S. (1973) Revised volcanic history of the San Juan, Uncompahgre, Silverton, and Lake City calderas in the western San Juan Mountains, Colorado: U.S. Geol. Survey Jour. Research, v. 1, no. 6, p. 627-642.
- Lipman, P. W., Steven, T. A., and Mehnert, H. H. (1970) Volcanic history of the San Juan Mountains, Colorado, as indicated by potassium-argon dating: *Geol. Soc. America Bull.*, v. 81, p. 2329-2352.
- Mehnert, H. H., Lipman, P. W., and Steven, T. A. (1973a) Age of the Lake City caldera and related Sunshine Peak Tuff, western San Juan Mountains, Colorado: *Isochron West*, v. 6, p. 31-33.
- _____ (1973b) Age of mineralization at Summitville, Colorado, as indicated by K-Ar dating of alunite: *Econ. Geol.*, v. 68, p. 399-401.
- Slack, J. F., and Lipman, P. W. (1979) Chronology of alteration, mineralization, and caldera evolution in the Lake City area, western San Juan Mountains, Colorado: *Nev. Bureau of Mines and Geol. Rept.* 33, p. 151-158.
- Steven, T. A., Cunningham, C. G., Naeser, C. W., and Mehnert, H. H. (1979) Revised stratigraphy and radiometric ages of volcanic rocks and mineral deposits in the Marysvale area, west-central Utah: U.S. Geol. Survey Bull. 1469.
- Steven, T. A., and Lipman, P. W. (1976) Calderas of the San Juan volcanic field, southwestern Colorado: U.S. Geol. Survey Prof. Paper 958, 35 p.