

## ***The Mule Mountain stock, an early Middle Devonian pluton in northern California***

J.P. Albers, R.W. Kistler, and L. Kwak

Isochron/West, Bulletin of Isotopic Geochronology, v. 31, pp. 17-18

Downloaded from: <https://geoinfo.nmt.edu/publications/periodicals/isochronwest/home.cfm?Issue=31>

---

Isochron/West was published at irregular intervals from 1971 to 1996. The journal was patterned after the journal *Radiocarbon* and covered isotopic age-dating (except carbon-14) on rocks and minerals from the Western Hemisphere. Initially, the geographic scope of papers was restricted to the western half of the United States, but was later expanded. The journal was sponsored and staffed by the New Mexico Bureau of Mines (now Geology) & Mineral Resources and the Nevada Bureau of Mines & Geology.



**ISOCHRON/WEST**  
*A Bulletin of Isotopic Geochronology*

All back-issue papers are available for free: <https://geoinfo.nmt.edu/publications/periodicals/isochronwest>

*This page is intentionally left blank to maintain order of facing pages.*

## THE MULE MOUNTAIN STOCK, AN EARLY MIDDLE DEVONIAN PLUTON IN NORTHERN CALIFORNIA

JOHN P. ALBERS  
RONALD W. KISTLER  
LORETTA KWAK

U.S. Geological Survey, Menlo Park, CA 94025 and Denver, CO 80225

The elliptical Mule Mountain stock west of Redding, Calif., is about 16 km long and 8 km wide and is composed mainly of highly altered and deeply weathered trondhjemite and albite granite. A very small area near its northern boundary is partly altered and unweathered hornblende quartz diorite. The stock intruded the Balaklala Rhyolite and Copley Greenstone of pre-late Middle Devonian age and is in turn intruded by the 134-m.y.-old Shasta Bally batholith. The age of the stock has been regarded on geologic grounds as Late Jurassic (Kinkel and others, 1956).

### DISCUSSION

A sample of quartz diorite has yielded concordant K-Ar and U-Pb ages. The concordance of the ages is remarkable in that even the  $^{208}\text{Pb}/^{232}\text{Th}$  age is within about 3 percent of the others, and the K-Ar age is about midway between the zircon extremes, well within their analytical uncertainty and eminently compatible. It is evident that the Mule Mountain stock is one of the few well-dated Paleozoic plutonic rocks in the western United States, and on the basis of the most recent estimate of the age of the Silurian-Devonian boundary (Ross and others, 1978), it appears that the stock is of early Middle Devonian age.

The pre-late Middle Devonian age of the Copley Greenstone and intertonguing Balaklala Rhyolite previously has been based on the fossiliferous Kennett Formation, which the Balaklala appears to underlie conformably. The new radiometric data not only support this age inference but also strongly suggest that volcanic rocks of the Copley and Balaklala, chiefly albitized andesite and dacite, may be generally cogenetic with Mule Mountain stock, as they are petrologically compatible. Barker and others (1979) concluded from petrologic studies that the trondhjemite of the Mule Mountain stock is cogenetic with extensive rocks of the Balaklala, and we conclude that inasmuch as the Copley Greenstone intertongues with the Balaklala, the Copley must also be coeval and cogenetic. The stock thus intruded its own volcanic pile during island arc magmatism in early Middle Devonian time. The stratabound massive sulfide deposits of the West Shasta mining district, which are enclosed within the Balaklala Rhyolite, were also in part a product of this event.

### SAMPLE DESCRIPTION

- K-Ar, U-Pb

Hornblende quartz diorite, unweathered and little altered (from "the very small area";  $40^{\circ}4'N$ ,  $122^{\circ}29'30''W$ ; W of Redding, CA). *Analytical data:* (Hornblende):  $\text{K}_2\text{O} = 0.182$  wt %,  $^*\text{Ar}^{40} = 11.483 \times 10^{-11}$  moles/gm,  $^*\text{Ar}^{40}/\Sigma\text{Ar}^{40} = 56\%$ ; (constants used)  $\text{K}^{40} = 0.01167$  atom %,  $\lambda_{\beta} = 4.962 \times 10^{-10}$  year $^{-1}$ ,  $\lambda_{\epsilon} + \lambda_{\epsilon}^1 = 0.581 \times 10^{-1}$  year $^{-1}$ . (Zircon): U = 520.3 ppm, Th = 196.7 ppm, Pb = 33.96 ppm;  $^{208}\text{Pb}/^{204}\text{Pb} = 5472.6$ ,  $^{207}\text{Pb}/^{204}\text{Pb} = 314.0$ ,  $^{208}\text{Pb}/^{204}\text{Pb} = 681.8$ .

(hornblende) K-Ar = 392  $\pm$  3 m.y.  
(zircon) Pb $^{208}$ /U $^{238}$  = 400 m.y.  
Pb $^{207}$ /U $^{235}$  = 400 m.y.  
Pb $^{208}$ /Pb $^{207}$  = 400 m.y.  
Pb $^{208}$ /Th $^{232}$  = 387 m.y.

[ISOCHRON/WEST, no. 31, August 1981]

### REFERENCES

- Barker, F., Millard, H. T., Jr., and Knight, R. J. (1979) Reconnaissance geochemistry of Devonian island-arc volcanic and intrusive rocks, West Shasta district, California: Trondhjemites, dacites and related rocks, Chapter 18 (Developments in Petrology) v. 6, p. 531.
- Kinkel, A. R., Jr., Hall, W. E., and Albers, J. P. (1956) Geology and base-metal deposits of the West Shasta copper-zinc district, Shasta County, California: U.S. Geological Survey Professional Paper 285.
- Ross, R. J., Jr., and others (1978) Devonian-Silurian boundary: U.S. Geological Survey open-file report 78-701, p. 363-365.

